

Bramido

FUNGICIDE

ACTIVE INGREDIENT	
Potassium sorbate.....	45%
INERT INGREDIENTS	
Water, Xantham Gum.....	55%
Total.....	100%

This product has not been registered by the U.S. EPA. Planta BioScience represents that this product qualifies for exemption from registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), Section 25b.

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.) **HAZARD & PRECAUTIONARY STATEMENTS:** Wear eye protection and chemical resistant protective gloves. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. **FIRST AID** Call a POISON CENTER or doctor/physician if you feel unwell. For Chemical Emergency (spill, leak, fire or accident), call CHEMTREC (800) 424-9300. **IF ON SKIN:** Take off immediately all contaminated clothing. Rinse skin with water/shower. Wash contaminated clothing before reuse. **IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. **IF INHALED:** Remove person to fresh air and keep comfortable for breathing. **IF SWALLOWED:** Rinse mouth. Do NOT induce vomiting.

GENERAL INFORMATION Bramido is a contact fungicide for field and greenhouse applications. Direct contact with the target organism is required for optimal control. Ensure application covers both sides of leaves. Bramido can be used in a disease resistance management program, rotating with fungicides having different modes of action. Bramido can also be used for late season disease infections when use of other fungicides may be limited due to pre-harvest interval restrictions.



MANUFACTURED BY
Planta BioScience
8696 Hinton Ave
Delhi CA 95315

Net Content: 2.5 gal
5 gal
275 gal

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS Avoid contact with eyes, skin and clothing. Harmful if swallowed.

Avoid breathing of spray mists. Avoid contamination of feed and foodstuffs. **STATEMENT OF PRACTICAL TREATMENT** If swallowed: Do not induce vomiting. Drink a glass of water and see a physician if experiencing gastrointestinal discomfort.

If on skin: Wash with soap and water. If in eyes:

Flush with water. If inhaled: Move to fresh air.

PERSONAL PROTECTIVE EQUIPMENT

APPLICATORS AND OTHER HANDLERS MUST WEAR: •Protective eyewear •Chemical resistant gloves •Longsleeved shirt and long pants •Shoes plus socks Follow manufacturer's instructions for cleaning/main- taining PPE. If no such instructions for washables, use detergent and hot water. **USER**

SAFETY RECOMMENDATIONS: Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash PPE separately from other laundry. **ENVIRONMENTAL HAZARDS** This product is for terrestrial uses only. Do not apply di-rectly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment wash-waters. Apply this product only as specified on this label.

STORAGE AND DISPOSAL 1. Prohibitions: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. 2. Storage: Store in a cool, dry, locked area out of the reach of children. Keep container tightly closed in storage to prevent entry of water. 3. Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. 4. Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

USE THIS PRODUCT IN A MANNER CONSISTENT WITH ITS LABELING DIRECTIONS FOR USE: Shake or Stir Concentrate to Uniform Consistency Prior to Opening. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Wait until sprays have dried before handling treated plants. Exempt from the requirements of the Worker Protection Standard, 40 CFR part 170.

FOR USE ON BUT NOT LIMITED TO: ornamental plants, orchard crops, berries, fruit trees, hops, hemp, vegetables, herbs, medicinal plants, cereals, pseudocereals, conifers, grasses, and alfalfa. **PESTS CONTROLLED:** Effective against fungal and bacterial

plant pathogens on a wide variety of crops. **USE PRECAUTIONS:** Keep the container tightly closed in storage. Do not spray plants under drought stress. Do not apply at temperatures exceeding 85°F. Do not tank mix with any herbicides or within spray tanks in which herbi- cides have been used. Application safety during the bloom period should be determined for each individual species of plant by conducting a small test. Do not tank mix with any herbicides or acidic products.

Maintain agitation through- out application. Only mix what can be sprayed at one time. Clean spray rig and lines following application. **MIXING INSTRUCTIONS:** All Phase must be mixed with water prior to application. Fill tank with $\frac{1}{2}$ to $\frac{3}{4}$ of the total required volume of water. Begin agitation of tank water and slowly add measured volume of All Phase. Continue agitating solution while adding remaining water. Agitate for at least ten minutes to fully activate ingredients prior to application.

APPLICATION INSTRUCTIONS: Use sufficient volume to completely cover all plant surfaces. **TIMING OF APPLICATIONS:** Applicator must adjust the precise timing and rate within the labeled range to fit local growth and climatic conditions.

DO NOT EXCEED MAXIMUM RATES OR APPLY MORE OFTEN THAN RECOMMENDED. MAY BE USED UP TO DAY OF HARVEST

APPLICATION RATES

In general, use the higher rates and/or shorter intervals to manage challenging diseases and/or when conditions are favorable for disease development. Complete coverage is essential for disease control.

Crops	Pest	RATES fl. oz./100 gal	COMMENTS
Almonds	Hull Rot (<i>Rhizopus stolonifer</i> , <i>Monilinia fructicola</i>)	32-128	Start application as soon as the first visual symptoms are observed and repeat every 7-10 days
Grapes	Powdery mildew (<i>Erysiphe necator</i>) Downy mildew (<i>Plasmopara viticola</i>) Sour rot (caused by a fungal complex) Botrytis (<i>Botrytis cinerea</i>) Black rot (<i>Guignardia bidwellii</i>) Phomopsis (<i>Phomopsis viticola</i>)	32-128	Start application as soon as the first visual symptoms are observed and repeat every 7-10 days. When conditions are favorable for downy mildew development during pre-flowering through fruit set stages. Ensure thorough coverage on the entire leaf surface.
Citrus	Greasy spot (<i>Mycosphaerella citri</i>)	32-128	Start application after petal fall or early fruit set and repeat every 21-28 days as long as conditions favor disease development.
Stone fruit, pome fruits and nuts	Powdery mildew (<i>Podosphaera</i> spp.) (<i>Erysiphe pistaciae</i>) (<i>Microsphaera penicillata</i>) Soft Fruit Rot	32-128	Start application as soon as the first visual symptoms are observed and repeat every 7-10 days. Ensure thorough coverage on the entire leaf surface.
Strawberry	Powdery mildew (<i>Podosphaera aphanis</i>) Botrytis (<i>Botrytis cinerea</i>)	32-128	Start application as soon as the first visual symptoms are observed and repeat every 7-10 days. Ensure thorough coverage on the entire leaf surface.
Blueberry	Botrytis blossom blight (<i>Botrytis cinerea</i>) Anthracnose fruit rot (<i>Collectotrichum acutatum</i>)	32-128	For Botrytis blossom blight, start a preventative application at early green tip and repeat every 7-10 days as long as conditions favor disease development. For Anthracnose fruit rot, start a preventative application at bloom and repeat every 7-10 days as long as conditions favor disease development.
Cucurbits	Powdery mildew (<i>Podosphaera xanthii</i>)	32-128	Start application as soon as the first visual symptoms are observed and repeat every 7-10 days. Ensure thorough coverage on the entire leaf surface.
Cannabis	Powdery mildew (<i>Podosphaera xanthii</i>)	32-128	Start application as soon as the first visual symptoms are observed and repeat every 7-10 days. Ensure thorough coverage on the entire leaf surface.